

Origin of the Neoproterozoic Baijianshan banded iron formation at the southeastern margin of the Tarim Block in NW China: implication for an extremely reducing ocean

Xiao-Shu Hao Chuan-Lin Zhang* Hui-Chao Zhang Teng Ding Xian-Tao Ye

College of Oceanography, Hohai University, Nanjing 210098, P.R. China

Abstract: The Neoproterozoic Banded iron formations (BIFs) were closely associated with the during the breakup of the Rodinia, thus they played an important role in our understanding of the atmospheric and oceanic oxygen levels during this period. In this contribution, the Neoproterozoic (ca. 737 Ma) Baijianshan BIF at Southeast Tarim, northwestern China was identified. Magnetite is the dominated iron-species, which occurs as the lamina interbedded with chert. The BIF contains low concentrations of trace elements, and is depleted in light rare earth elements (LREEs) based on comparison with the Post Archean Australian Shale (PAAS). In addition, the BIF exhibits slightly positive La-Eu anomalies, negligible Ce anomalies, insignificant Y anomalies, chondritic Y/Ho ratios (23-32), and slightly chondritic initial $^{57}\text{Fe}_{\text{IRMM-014}}$ values ranging from -0.45 to 1.46, averaging 0.37). All these features indicate that the precipitation of Baijianshan BIF was closely related to the submarine low-T hydrothermal fluids with little detrital contribution. Moreover, the Baijianshan BIF is revealing the partial oxidation of Fe^{2+} into Fe^{3+} during the precipitation of this BIF. Our data suggest that the formation of Baijianshan BIF was closely associated with a significantly reducing ocean, which most likely was isolated from the oxidized atmosphere by a local ice sheet. This Neoproterozoic Baijianshan ocean has the initial oxygen levels as low as, or even lower than that of Archean and Paleoproterozoic oceans.

Key words: Neoproterozoic, banded iron formations (BIFs), Sr-Nd isotope, Fe isotope, Tarim block

*Corresponding Author: E-Mail: zhangchuanlin@hhu.edu.cn

1 Introduction

Banded iron formations (BIFs), one of the most important iron sources in the world, are typical marine sedimentary rocks mainly precipitated during the Precambrian. BIF was firstly defined as a type of sedimentary rock containing more than 15% iron (James, 1954, 1966), which was later revised as thinly layered or laminated rock in which chert (or its metamorphic equivalent) alternating with layers that are composed mainly of iron minerals with the iron content ranging from 20% to 35% (James, 1983). The oldest BIFs precipitated at 3.8 Ga (Isua, Greenland), and then a large number of BIFs formed during the Archean-Paleoproterozoic (3.5-1.8 Ga), with a peak at ca. 2.5 Ga (Klein, 2005 and the references therein). They disappeared at ca. 1.8 Ga, and reoccurred in the Neoproterozoic (0.8-0.6 Ga) after a billion-year hiatus (Klein, 2005). It is commonly accepted that the precipitations of BIFs were closely related to fundamental changes in atmospheric and/or oceanic oxygen levels (i.e., Great Oxidation Event, GOE), although some key issues, including the source of the hydrothermal fluids, oxygen level and tectonic settings, are still under debate (Holland, 2002). In line with tectonic backgrounds, the BIFs can be subdivided into Algoma- and Superior-type (Gross, 1980), the Algoma-type generally precipitated nearby the volcano craters, in the sequences mainly composed of magnetite and hematite (Klein and Beukes, 1992; Li et al., 2014; Sial et al., 2015), and the Superior-type precipitated in shallow marine environment, at passive continental margin (Gross and McLeod, 1980).

Although the resources of Neoproterozoic BIFs are even much less than those of Archean and Paleoproterozoic, they are widely distributed worldwide, such as Rapitan BIF in Canada (Halverson et al., 2002) and Um Anab BIF in Egypt (Basta et al., 2011), Sawawin BIF in Saudi Arabia (Freitas et al., 2011), Holowilena and Oraparinna in South Africa (Freitas et al., 2011), Boston Peak Formation in California (Lechte et al., 2018) and Wadi el-Hadid in the Arabian Shield (El-Rahman et al., 2020). They were also discovered in the Neoproterozoic orogenic belts of South China, such as Sanjiang in Guangxi (Li et al., 2018), Shilu in Hainan Province (Xu et al., 2014), Xinyu in Jiangxi Province (Li et al., 2018; Zhu et al., 2019), the Shalong in the Central Tianshan (Lei et al., 2016, 2017), and the Watan BIF in the Western Kunlun orogenic belt (Hu et al., 2017, 2020). Several

Conventional magnetic and density techniques were used to concentrate non-magnetic, heavy fractions and then zircon grains were hand-picked under a binocular microscope. After being mounted in epoxy mount, zircon grains were then polished to section the crystals in half for analysis. All zircons were documented with transmitted and reflected light micrographs as well as cathodoluminescence (CL) images to reveal their internal structures. Zircon U-Pb ages were analyzed using the LA-ICP-MS at the Tianjin Institute of Geology and Mineral Resources (China Geological Survey). Detailed analytical procedures can be found in Li et al. (2009). Zircon standards 91500 and GJ-1 (Jackson et al., 2004) were used to monitor the analysis. The U-Pb concordia plots were processed by ISOPLOT 3.0 and data were presented with 1 σ errors and 95% confidence limits (Ludwig, 2003). The zircon U-Pb age data are listed in Supplementary Table 1.

4.2 Whole rock geochemistry

Seventeen BIF samples and three basalt samples were carefully selected along the geological location. Whole-rock major compositions were analyzed using standard X-ray fluorescence (XRF) on fused glass beads at the Nanjing Institute of Geology and Mineral Resources (China Geological Survey), following the procedures similar to those described by Li et al., (2006a, b). Analytical uncertainties are between 1 and 5%. In addition, we used wet chemistry technique to measure FeO and Fe₂O₃, with the procedures described by Andrade et al. (2002). The measured data are listed in supplementary Table 2.

Trace elements were determined using a Perkin-Elmer Sciex ELAN DRC-e ICP-MS at the State Key Laboratory of Ore Deposit Geochemistry, Institute of Geochemistry (Chinese Academy of Sciences), with the analytical accuracy better than 5%. Samples were digested with 1ml of HF and 0.5 ml of HNO₃ in screw top PTFE-lined stainless steel bombs at 190 $^{\circ}$ C for 12h (Qi et al., 2000). The analytical precision for most elements was better than 1% with concentrations > 200 ppm, and 1-3% when less than 200 ppm. The analytical results are reported in supplementary Table 2.

Sr-Nd isotopes were measured using the Micromass Isoprobe Multi-collector MC-ICP-MS at Tianjin Institute of Geology and Mineral Resources (China Geological Survey), with the analytical procedures similar to those reported by Li et al. (2004). Measured ⁸⁷Sr/⁸⁶Sr and ¹⁴³Nd/¹⁴⁴Nd ratios were corrected for mass-fractionation using ⁸⁶Sr/⁸⁸Sr = 0.1194 and ¹⁴⁶Nd/¹⁴⁴Nd = 0.7219, respectively. The measured values for the NBS SRM 987 standard ⁸⁷Sr/⁸⁶Sr = 0.71025 and the Shin Etsu JNdi-1 standard ¹⁴³Nd/¹⁴⁴Nd = 0.512115. All the initial ¹⁴³Nd/¹⁴⁴Nd values of the BIF are calculated with a probable depositional age (737 Ma) for the Baijianshan BIF.

Iron isotope ratios were measured using MC-ICP-MS at Beijing Createch Testing Technology Co., Ltd. The results of Fe isotope ratio determination as the ten thousandth deviation of the sample from the standard sample is as follows:

$$\delta^x Fe/^{54}Fe(\text{‰}) = \left[\frac{(^x Fe/^{54} Fe)_{sample}}{(^x Fe/^{54} Fe)_{standard}} - 1 \right] \times 1000$$

(x = 56, 57)

The performance of the instrument was assessed by repeated measurements of an in-house standard (CAGS Fe) that yielded deviations relative to the IRMM-014 Fe isotope reference material. The

$$^{56}Fe_{CAGS} \quad ^{56}Fe_{IRMM014} \quad ^{56}Fe_{CAGS} \quad ^{57}Fe_{IRMM014}$$

ribed by Tang et al. (2016).

the significant $^{206}\text{Pb}/^{238}\text{U}$ age, yield a mean $^{206}\text{Pb}/^{238}\text{U}$ age of 737.0 ± 4.0 Ma (N=15, MSWD=0.9), which is interpreted as the deposition time of the Baijianshan BIF.

Whole rock Geochemistry
Geochemistry of the basalts

The basalt samples (2702H1, 270H2 and 2702H3) have low SiO_2 and high CaO contents. The trace element patterns (compatible elements for the rock type classification are used) show a typical alkaline signature with La_N/Yb_N ratios (1.05-1.09) define their alkaline signature and in the field of the alkaline basalt field (Figure not shown). With respect to REE patterns, the samples show a typical alkaline signature ranging from 186 ppm to 195 ppm and show variable La_N/Yb_N ratios (Figure 5a) ($\text{La}_N/\text{Yb}_N = 9.8-11.0$). In line with their low La_N/Yb_N ratios, the basalts are evolved magma. However, the patterns of the OIB-like basalts with insignificant enrichment of the

Geochemistry of the BIF

shown in supplementary Table S1. The REE patterns and concentrations vary between samples. The SiO_2 content is 26.81% and 55.81% (Table S1). The CaO content is 1.56% and 1.56% (Table S1). The trace element patterns of the BIF samples are similar to those of the basalts.

The BIF samples show a typical alkaline signature with La_N/Yb_N ratios (1.05-1.09) define their alkaline signature and in the field of the alkaline basalt field (Figure not shown). With respect to REE patterns, the samples show a typical alkaline signature ranging from 186 ppm to 195 ppm and show variable La_N/Yb_N ratios (Figure 5a) ($\text{La}_N/\text{Yb}_N = 9.8-11.0$). In line with their low La_N/Yb_N ratios, the basalts are evolved magma. However, the patterns of the OIB-like basalts with insignificant enrichment of the

The Baijianshan BIF range from 0.02 to 3.47. The $\text{Fe}_{\text{IRMM-014}}$ values of 1.24-1.47. The Fe isotopes shares the signature of the BIF (Shen et al., 2011; Shen et al., 2018).

Resources for the Baijianshan BIFs

and microscopic observations

experienced significant metamorphism (Fig. 3b, c and d), indicating the original geochemical characteristics were well retained and can be used to decipher the iron sources and the oceanic environments when it precipitated.

The Baijianshan BIF shows a modern seawater REE signature with significant LREE depletion relative to the HREE ($La_N/Yb_N = 0.41$) (Fig. 5b, Zhang and Nozaki, 1996; Alibo and Nozaki, 1999). A range of factors can affect the primary REY composition in terms of BIF depositional processes, including post-depositional metamorphism and/or syn-deposition clastic contamination (Alexander et al., 2008; Hu et al., 2020). Previous studies demonstrated that diagenetic and metamorphic events (such as weathering and/or fluid-rock interactions) have negligible effects on the REY of the chemical sediments (Bolhar et al., 2004). However, Rb is a relatively mobile element compared to highly immobile elements such as Th. In Fig. 6a, the well positive correlation between Rb and Th demonstrates insignificant mobility of the LILEs (Large Ion Lithophile Elements) and the absence of significant diagenetic or metamorphic alteration effects, this conclusion is also consistent with thin section observations (Fig. 3c, d, g, h). With respect to the syn-depositional processes, the contents of Al_2O_3 and TiO_2 are consistently low with averages of 3.66% and 0.41%. In addition, despite the positive correlations between Al_2O_3 and TiO_2 ($r = 0.93$), K_2O ($r = 0.79$), Rb ($r = 0.78$), V ($r = 0.45$), Sc REE ($r = 0.62$) (Fig. 6b-i) signify the incorporation of terrigenous contribution in the deposition processes, most correlation coefficients are much lower than those BIFs sourced from recycled crust (Basta et al., 2011; Cox et al., 2013; Hu et al., 2017, 2020). Besides, the Baijianshan BIF shares similar REE patterns with those BIFs from North China Craton but contains slightly higher contents of Al_2O_3 and TiO_2 (Li et al., 2014). Taken together, we suggest a little incorporation of terrigenous component in the Baijianshan BIF.

Positive Eu anomalies are generally interpreted as the precipitation of high-temperature hydrothermal fluids (Danielson et al., 1992; Bau and Dulski, 1999) while low-temperature hydrothermal fluids usually display weak or no Eu anomalies (Michard et al., 1993; Li et al., 2014). Most samples from the Baijianshan BIF show negligible Eu anomalies ($Eu/Eu^* = 0.96\sim 1.14$, averaging 1.07), which are indicative of low-temperature hydrothermal solutions (Danielson et al., 1992). The

(t) value can be used as a tracer to distinguish different sources and crustal contamination. As shown in Fig. 7, the Baijianshan BIF has slightly average (t) value (average of 0.37), much lower than those of Baijianshan basalt, indicating that the deposition of the Baijianshan BIF was controlled by a hydrothermal flux with mantle- (Jacobsen and Pimentel-Klose, 1988). However, the slightly negative correlation between (Ma) and Al_2O_3 contents (Fig. 8a) argues for the low proportion of detrital components inputting.

The Y/Ho ratios of BIFs can provide constraints of the material source and precipitation environment of BIFs (Hu et al., 2017). Modern seawaters have a Y/Ho ratio of 44-74 (Bau and Dulski, 1996). The terrestrial material has a Y/Ho ratio of ~26, any little terrestrial contamination could quickly descend seawater-like superchondritic Y/Ho ratios (>44) (Bolhar et al., 2004). The hydrothermal fluids have almost chondritic Y/Ho ratios (26-28, Douville et al., 1999; Bau and Dulski, 1999). The Baijianshan BIF has a range of Y/Ho ratios between 23 and 32, similar to the chondritic values (26-28, Bau and Dulski, 1999), possibly due to the mixture of high and/or low-T hydrothermal fluids (Hu et al., 2020). Y/Ho ratios of the Baijianshan BIF samples might inherit from the low-T hydrothermal fluids, but the influences of terrestrial materials can not be completely excluded. Compared with the BIFs in North China Craton, the Baijianshan BIF displays relatively low positive Eu anomalies, low Y/Ho ratios as well as a slightly depleted (Li et al., 2014; Wang et al., 2016), illustrating the input of low-temperature hydrothermal fluids during the deposition of the Baijianshan BIF.

Ce/Ce*_{PAAAS} values with variable Al₂O₃ contents of the Baijianshan BIF indicate that terrigenous detrital materials played little role in the geochemical budget of Ce. In oxidized seawater, Ce(III) transforms into Ce(IV), and then Ce(IV) is likely to be hydrolyzed and precipitated with Fe-Mn oxyhydroxides, organics and clay which lead to a significant negative Ce anomaly of seawater (Byrne and Sholkovitz, 1996). As shown in the Ce/Ce* vs. Pr/Pr* diagram (Fig. 10), all the Baijianshan BIF samples display no Ce anomalies, similar with most Neoproterozoic BIFs, unambiguously arguing for anoxic environment in the ancient ocean. Although the atmospheric and oceanic oxygen levels during the Neoproterozoic are still under debate, numerous studies revealed that Neoproterozoic oceans underwent a stepwise and protracted oxidation. Anoxic ferruginous deep seawater was a typical feature of the late Neoproterozoic, as inferred from geochemical proxies such as iron geochemistry (e.g., Canfield et al., 2008; Sperling et al., 2015), redox-sensitive elements (e.g., Schröder and Grotzinger, 2007; Rajabi et al., 2015), and framboidal pyrite (e.g., Rajabi et al., 2015). Stern et al. (2013) suggested that this scenario might be attributed to the seawater surfaces being covered by ice sheets, blocking the oxygen from atmosphere dissolving into the widespread anoxic ferruginous ocean. Despite no tillite was identified at the Baijianshan BIF, the coeval tillite sequences were documented in the Quruqtagh area (i.e., ca.740 Ma Beiyixi tillite) of NE Tarim (Xu et al., 2009), which argued for the possibility of ice sheets in the Cryogenian Baijianshan ocean.

In a Fe²⁺-rich marine settings, both abiotic and biotic action were demonstrated as oxidation pathways for the precipitation of iron oxyhydroxides (Fe(OH)₃) from dissolved Fe²⁺ (Konhauser et al., 2011). As an element of variable valency, the fractionation of Fe isotope is affected by the oxidation-reduction condition (Bullen et al., 2001; Zhu et al., 2002; Balci et al., 2006). Experimental studies demonstrated that oxidation of Fe²⁺ to Fe³⁺ in solution causes considerable Fe isotope fraction, generating an enrichment of heavy Fe isotopes in Fe³⁺ (Bullen et al., 2001; Johnson et al., 2002; Balci et al., 2006; Hou et al., 2014). On the other hand, the fractionation caused by pyrite with light-Fe isotopes is considered to play a leading role in the enrichment of heavy Fe isotopes in the residual system relative to the original state (Rouxel et al., 2016), since the pyrite is easier to precipitation from plume. Nevertheless, there is no sulfide observed in the Baijianshan BIF, thus the potential influences of sulfides on the enrichment of heavy Fe isotopes can be excluded. Under oxidized environment, the enrichment of heavy Fe isotopes is shown in oxide or hydroxide of iron trivalent (Fe³⁺_{ppt}) and light Fe isotope is displayed in Ferrous solution (Fe²⁺_{aq}) (Johnson et al., 2002). The magnitude of Fe isotope fractionation is controlled by the degree of precipitation from Fe²⁺ to Fe³⁺, which is related to the degree of ocean oxidation (Yan et al., 2010). The Fe in the seawater can be completely precipitated and no fractionation of the Fe isotope occurs when the seawater is completely oxidized, thus the Fe isotope values in iron oxide precipitates (Fe³⁺) can represent the Fe isotope information of seawater. However, when the seawater is partially oxidized, the Fe in the seawater is partially precipitated and the Fe isotope fractionation will occur in iron oxide precipitates (Fe³⁺). Therefore, the Fe isotope value can be used as a proxy to decipher the redox state of seawater (Yan et al., 2010). Compared with some BIFs deposited during the Archean-Paleoproterozoic (Dauphas et al., 2004; Rouxel et al., 2005; Li et al., 2012; Johnson et al., 2003; Hou et al., 2014; Li et al., 2014), the Neoproterozoic Baijianshan Sanjiang (Yan et al., 2010) and Xinyu BIFs (Shen et al., 2008) have higher values ⁵⁷Fe (Fig. 11). Several studies demonstrated that the low ⁵⁷Fe values of BIFs could be due to the contributions from a continental component and the continental Fe source is best explained by Fe mobilization on the continental margin by microbial dissimilatory iron reduction (DIR) (Johnson et al., 2008; Li et al., 2015). As for the Fe isotope values of Archean-Paleoproterozoic BIFs we cited, the Fe source of SW Greenland BIF, Anshan-Benxi BIFs, Gongchangling BIFs and Yuanjiacun BIFs are unambiguously demonstrated to be hydrothermal type similar to the scenario of the Baijianshan BIF (Dauphas et al., 2004; Li et al., 2012; Li et al., 2014; Hou et al., 2014), while the Fe of Transvaal, Manjeri, Belingwe Belt and Zimbabwe BIFs are probably from multiple sources (abiologic and biologic processes) (Johnson et al., 2008; Grassineau et al., 2011). The comparison reveals that the environment in which they precipitated was more reduced. Both the iron isotope and elemental geochemistry thus demonstrate that the Neoproterozoic Baijianshan ocean might have the same reduced environment as Archean or Paleoproterozoic, or even more reduced.

6.3 A model of the Baijianshan BIF

During the late Neoproterozoic period (i.e., 750-550 Ma), the Earth experienced long-lived global glaciations, known as the Snowball Earth events (Hoffman et al., 1998). In view of the Neoproterozoic oxygenation event (NOE) at that time, the content of oxygen in atmosphere was much higher than before (Och et al., 2012). Recently, EI-Rahman et al. (2020) argued that some Neoproterozoic BIFs are not glaciogenic due to their significant depleted Nd isotope compositions. Under this scenario, the iron was most likely deriving from hydrothermal alteration of juvenile oceanic crust. The Chondritic-like Nd isotope compositions of the Baijianshan BIF (do not favor this model. Furthermore, the close temporally-spatially related tillite sequences in the Quruqtagh of NE Tarim (Xu et al., 2009), 765 Ma-735 Ma tillites in Namibia, suggested a local glaciation termed as Kaigas

glaciation (Kay et al., 2001; Cailteux et al., 2005; Hoffmann et al., 2006). Considering the high latitude position of the Tarim in the Rodinia configuration (Li et al., 2008), the Kaigas glaciation (corresponding to the Beiyixi tillite in Tarim) most possibly occurred in Tarim. In combination with previous studies on other BIFs with the data presented in this study, we construct a cartoon model showing the formation mechanism of the Baijianshan BIF (Fig. 12).

Stage 1, the seawater was covered by ice sheet which led to the insulation between atmosphere and oceans with the in

And then it induced the emergence of a reductive environment. The iron supplied by low-temperature hydrothermal fluids could exist as the ferrous form, forming a relatively stable anoxic ferruginous reservoir in the ocean.

Stage 2, at interglacial period, the partial melting of the ice sheet caused some oxygen from the atmosphere into the ocean, which emerged a stratified seawater. There were mainly anoxic and Fe²⁺-rich deeper seawater, whereas the Fe²⁺ was gradually oxidized into Fe³⁺ and precipitated into iron oxide in near-shore oxic shallow seawaters. The low-temperature hydrothermal was the dominant source during the precipitation process to interpret the slight positive Eu anomalies and positive values

. Under the effect of the melting ice-sheet, the increasing weathering then contributed to a small amount of terrestrial materials into the shallow seawater, resulting the precipitation of the Baijianshan BIF.

7 Conclusions

The Baijianshan BIF precipitated in glacially-influenced settings during the Neoproterozoic (737 Ma). Low-T hydrothermal fluids, accompanied with limited detritus input, were the dominated iron sources for the formation of Baijianshan BIF.

The Baijianshan ocean was initially covered by ice sheets, giving rise to the extremely reduced environment favored by Fe²⁺. Then it was oxidized into Fe³⁺ and precipitated as magnetite due to the thawing of ice sheets. At the early stage, local reduced ocean enhanced Fe fluxes genetically related to the intensive marine magmatism and hydrothermal activities due to the existence of ice sheets. Thawing of ice sheets induced oxidized oceanic layer, favoring the formation of the Neoproterozoic BIF at the later stage.

Acknowledgements

We sincerely thank the anonymous reviewers for their careful and constructive comments on our manuscript, which significantly elevate the quality of this study. We are grateful to Prof. Hai-Bo Zou, for his careful polishing on the English writing. This project is funded by the Fundamental Research Fund for Central Universities (B16020127).

References

- Alexander, B.W., Bau, M., Andersson, P., Dulski, P., 2008. Continentally-derived solutes in shallow Archean seawater: rare earth element and Nd isotope evidence in iron formation from the 2.9 Ga Pongola Supergroup, South Africa. *Geochim. Cosmochim. Acta* 72, 378-394.
- Alibo, D.S., Nozaki, Y., 1999. Rare earth elements in seawater: particle association, shale normalization and Ce oxidation. *Geochim. Cosmochim. Acta* 63(3-4), 363-372.
- Andrade, S., Hypolito, R., Ulbrich, H.H.G.J., Silva, M.L., 2002. Iron(II) oxide determination in rocks and minerals. *Chem. Geol.* 182, 85-89.
- Balci, N., Bullen, T.D., Witte-Lien, K., Shanks, W.C., Motelica, M., Mandernack, K.W., 2006. Iron isotope fractionation during microbially stimulated Fe(II) oxidation and Fe(III) precipitation. *Geochim Cosmochim Acta* 70, 622-639.
- Basta, F.F., Maurice, A.E., Fontboté, L., Favarger, P.Y., 2011. Petrology and geochemistry of the banded iron formation (BIF) of Wadi Karim and UmAnab, Eastern Desert, Egypt: implications for the origin of Neoproterozoic BIF. *Precambrian Res.* 187, 277-292.
- Bau, M., Dulski, P., 1996. Distribution of yttrium and rare-earth elements in the Penge and Kuruman iron-formations, Transvaal Super group, South Africa. *Precambrian Res.* 79, 37-55.
- Bau, M., Dulski, P., 1999. Comparing yttrium and rare earths in hydrothermal fluids from the Mid-Atlantic Ridge: implications for Y and REE behaviour during nearvent mixing and for the Y/Ho ratio of Proterozoic seawater. *Chem. Geol.* 155, 77-90.
- Bolhar, R., Kamber, B.S., Moorbath, S., Fedo, C.M., Whitehouse, M.J., 2004. Characterisation of early Archaean chemical sediments by trace element signatures. *Earth Planet. Sci. Lett.* 222, 43-60.
- Bullen, T.D., White, A.F., Childs, C.W., Vivit, D.V., Schulz, M.S., 2001. Demonstration of significant abiotic iron isotope fractionation. *Geology* 29, 699-702.
- Byrne, R., Sholkovitz, 1996. Marine chemistry and geochemistry of the lanthanides. In: Gschneider Jr.K.A., Eyring, L. (Eds.), *Handbook on the Physics and Chemistry of the Rare Earths*. Elsevier, Amsterdam.
- Cailteux, J.L.H., Kampunzu, A.B.H., Batumike, M.J., 2005. Lithostratigraphic position and petrographic Journal of African Earth Sciences 42, 82-94.
- Canfield, D.E., Poulton, S.W., Knoll, A.H., Narbonne, G.M., Ross, G., Goldberg, T., Strauss, H., 2008. Ferruginous Conditions Dominated Later Neoproterozoic Deep-Water Chemistry. *Science* 321(5891), 949-952.
- Chen, D.Z., Wang, J.G., Qing, H.R., Yan, D.T., Li, R.W., 2009. Hydrothermal venting activities in the Early

- Cambrian, South China: Petrological, geochronological and stable isotopic constraints. *Chemical Geology* 258, 168-181.
- Cox, G.M., Halverson, G.P., Minarik, W.G., Heron, D.P.L., Macdonald, F.A., Bellefroid, E.J., Strauss, J.V., 2013. Neoproterozoic iron formation: an evaluation of its temporal, environmental and tectonic significance. *Chem. Geol.* 362, 232-249.
- Cox, G.M., Halverson, G.P., Poirier, A., Le Heron, D., Strauss, J.V., Stevenson, R., 2016. A model for Cryogenian iron formation. *Earth Planet. Sci. Lett.* 433, 280-292.
- Danielson, A., Moller, P., Dulski, P., 1992. The europium anomalies in banded iron formations and the thermal history of the oceanic crust. *Chem. Geol.* 97, 89-100.
- Dauphas, N., van Zuilen, M., Wadhwa, M., Davis, A.M., Marty, B., Janney, P.E., 2004. Clues from Fe isotope variations on the origin of early Archean BIFs from Greenland. *Science* 306, 2077-2080.
- EI-Rahman, Y.A., Gutzmer, J., Li, X.H., Seifert, T., Li, C.F., Ling, X.X., Li, J., 2020. Not all Neoproterozoic iron formations are glaciogenic: Sturtian-aged non-Rapitan exhalative iron formations from the Arabian-Nubian Shield. *Mineralium Deposita* 55, 577-596.
- Feng, L.J., Huang, J., Lu, D.B., Zhang, Q.R., 2016. Major and trace element geochemistry of the Neoproterozoic syn-glacial Fulu iron formation, South China. *Geological Magazine* 154, 1-10.
- Freitas, B.T., Warren, L.V., Boggiani, P.C., Almeida, R.P.D., Piacentini, T., 2011. Tectono-sedimentary evolution of the Neoproterozoic BIF-bearing Jacadigo Group, SW-Brazil. *Sedimentary Geology* 238, 48-70.
- Grassineau, N.V., Nisbet, E.G., Bickle, M.J., Fowler, C.M.R., Lowry, D., Mathey, D.P., Abell, P., Martin, A., 2000. Antiquity of the biological sulphur cycle: evidence from sulphur and carbon isotopes in 2700 million-year-old rocks of the Belingwe Belt, Zimbabwe. *Proc. R. Soc. Lond. B* 268, 113-119.
- Gross, G.A., 1980. A classification of iron-formation based on depositional environments. *Canadian Mineralogist* 18, 215-222.
- Gross, G.A., Mcleod, C.R., 1980. A preliminary assessment of the chemical composition of iron formation in Canada. *Canadian Mineralogist* 18, 223-229.
- Halverson, G.P., Poitrasson, F., Hoffman, P.F., Nédélec, A., Montel, J.M., Kirby, J., 2011. Fe isotope and trace element geochemistry of the Neoproterozoic syn-glacial Rapitan iron formation. *Earth Planet. Sci. Lett.* 309(1), 100-112.
- Hoffman, P.F., Kaufman, A.J., Halverson, G.P., Schrag, D.P., 1998. A Neoproterozoic snowball Earth. *Science* 281, 1342-1346.
- Hoffman, P.F., Schrag, D.P., 2002. The snowball earth hypothesis: testing the limits of global change. *Terra Nova* 14(3), 129-155.
- Hoffmann, K.H., Condon, D.J., Bowring, S.A., Prave, A.R., Fallick, A., 2006. Lithostratigraphic, carbon (^{13}C) isotope and U-Pb zircon age constraints on early Neoproterozoic (ca. 755 Ma) glaciation in the Gariep Belt, southern Namibia. *Snowball Earth Conference*, July 16-21, 2006, Montle Namibia, Ticino, Switzerland, 51.
- Holland, H.D., 2002. Volcanic gases, black smokers, and the great oxidation event. *Geochimica et Cosmochimica Acta* 66(21), 3811-3826.
- Hou, K.J., Li, Y.H., Gao, J.F., Liu, F., Qin, Y., 2014. Geochemistry and Si-O-Fe isotope constraint on the origin of banded iron formations of the Yuanjiacun Formation, Lvliang Group, Shanxi, China. *Ore Geology Reviews* 57, 288-298.
- Hu, J., Wang, H., Wang, M., 2017. Geochemistry and origin of the Neoproterozoic Dahongliutan banded iron formation (BIF) in the Western Kunlun orogenic belt, Xinjiang (NW China). *Ore Geology Reviews* 89, 836-857.
- Hu, J., Wang, H., Zhang, L.G., 2020. A rare earth element and Nd isotopic investigation into the provenance and deposition of the Dahongliutan banded iron formation and associated carbonates, NW China: Implications on Neoproterozoic seawater compositions. *Precambrian Research* 342, 1-29.
- Jackson, S.E., Pearson, N.J., Griffin, W.L., Belousova, E.A., 2004. The application of laser ablation-inductively coupled plasma-mass spectrometry to in situ U Pb zircon geochronology. *Chemical Geology* 211(1-2), 47-69.
- Jacobsen, S.B., Pimentel-Klose, M.R., 1988. Nd isotopic variations in Precambrian banded iron formations. *Geophys Res Lett* 15, 393-396.
- James, H.L., 1954. Sedimentary facies of iron-formation. *Economic Geology* 49(3), 235-293.
- James, H.L., 1966. Chemistry of the iron-rich sedimentary rocks. *U.S. Geol. Survey Prof.* 440-461.
- James, H.L., 1983. Distribution of banded iron-formation in space and time. *Developments in Precambrian Geology* 6, 471-490.
- Johnson, C.M., Beard, B.L., Beukes, N.J., 2008. Earth as inferred from Fe isotope studies of banded iron formations from the Transvaal Craton. *Contributions to Mineralogy and Petrology* 144(5), 523-547.
- Johnson, C.M., Beard, B.L., Klein, C., Beukes, N.J., Roden, E.E., 2008. Iron isotopes constrain biologic and abiologic processes in banded iron formation genesis. *Geochimica Et Cosmochimica Acta* 72(1), 151-169.
- Johnson, C.M., Skulan, J.L., Beard, B.L., Sun, H., Neelson, K.H., Braterman, P.S., 2002. Isotopic fractionation between Fe(III) and Fe(II) in aqueous solution. *Earth Planet Sci Lett* 195, 141-153.
- Johnston, D.T., Poulton, S.W., Dehler, C., Porter, S., Husson, J., Canfield, D.E., Knoll, A.H., 2010. An emerging picture of Neoproterozoic ocean chemistry: insights from the Chuar Group, Grand Canyon, USA. *Earth Planet. Sci. Lett.* 290, 64-73.
- Kay, R.M., Liyungu, A.K., Njamu, F.M., Somwe, V., Banda, J., Mosley, P.N., Armstrong, R.A., 2001. The western arm of the Lufilian Arc in NW Zambia and its potential for copper mineralization. *Journal of African Earth Sciences* 33, 503-528.
- Klein, C., 2005. Some Precambrian banded iron-formations (BIFs) from around the world: Their age, geologic setting, mineralogy, metamorphism, geochemistry, and origins. *American Mineralogist* 90(10), 1473-1499.
- Klein, C., Beukes, N.J., 1992. Time distribution, stratigraphy, sedimentologic setting, and geochemistry of Precambrian iron-formations. In: Schopf, J.W., Klein, C. (Eds.), *The Proterozoic Biosphere*. Cambridge University Press, Cambridge, 139-146.
- Klein, C., Ladeira, E.A., 2004. Geochemistry and mineralogy of Neoproterozoic banded iron-formations and some selected, siliceous manganese formations from the Urucum district, Mato Grosso do Sul, Brazil. *Econ. Geol.* 99, 1233-1244.
- Konhauser, K.O., Kappler, A.K., Roden, E.E., 2011. The microbial role in iron redox and biomineralization reactions. *Elements* 7, 89-93.

- change. *Precambr. Res.* 239, 79-94.
- Sun, S.S., McDonough, W.F., 1989. Chemical and isotopic systematics of oceanic basalts: Implications for mantle composition and processes. *Geological Society London Special Publications* 42(1), 313-345.
- Tang, S.H., Zhu, X.K., Li, J., Yan, B., Li, S.Z., Li, Z.H., Wang, Y., Sun, J., 2016. New Standard Solutions for Measurement of Iron, Copper and Zinc Isotope Compositions by Multi-collector Inductively Coupled Plasma-Mass Spectrometry. *ROCK AND MINERAL ANALYSIS* 35(2), 127-133 (in Chinese with English abstract).
- Wu, Y.B., Zheng, Y.F., 2004. A study on the genetic mineralogy of zircon and its restriction on the interpretation of U-Pb age. *CHINESE SCIENCE BULLETIN* 49(16), 1589 (in Chinese).
- Xinjiang Bureau of Geology and Mineral Resources (BGMR), 1993. *Regional Geology of Xinjiang Uygur Autonomous Region*. Geological Publishing House, Beijing, 8-33 (in Chinese).
- Xu, B., Xiao, S.H., Zou, H.B., Chen, Y., Li, Z.X., Song, B., Liu, D.Y., Zhou, C.M., Yuan, X.L., 2009. SHRIMP zircon U-Pb age constraints on Neoproterozoic Quruqtagh diamictites in NW China. *Precambrian Research* 168, 247-258.
- Xu, B., Zou, H.B., Chen, Y., He, J.Y., Wang, Y., 2013. The Sugetbrak basalts from northwestern Tarim Block of northwest China: Geochronology, geochemistry and implications for Rodinia breakup and ice age in the Late Neoproterozoic. *Precambrian Research* 236, 214-226.
- Xu, D.R., Wang, Z.L., Chen, H.Y., Hollings, Pete, Jansen, Nicholas, H., Zhang, Z.C., Wu, C.J., 2014. Petrography and geochemistry of the Shilu Fe Co Cu ore district, South China: Implications for the origin of a Neoproterozoic BIF system. *Ore Geology Reviews* 57, 322-350.
- Xu, Z.Q., He, B.Z., Zhang, C.L., Zhang, J.X., 2013. Tectonic framework and crustal evolution of the Precambrian basement of the Tarim Block in NW China: New geochronological evidence from deep drilling samples. *Precambrian Research* 235, 150-162.
- Yan, B., Zhu, X.K., Tang, S.H., Zhu, M.Y., 2010. Fe isotopic characteristics of the Neoproterozoic BIF in Guangxi Province and its Implications. *ACTA GEOLOGICA SINICA* 33(4), 761-766 (in Chinese with English abstract).

Figure captions




Fig. 1 (a) Sketch Precambrian geological map of Tarim Craton, showing the location of the Baijianshan BIF (modified after Lu et al., 2008). (b) Geological map of the Baijianshan Banded Iron Formation (Baijianshan BIF).




Fig. 2 Stratigraphical column of the Baijianshan BIF.




Fig. 3 Field photographs, hand specimen and mineral features of the Baijianshan BIF. (a) The tuff layers in the Baijianshan BIF. (b) Laminated Fe orebodies with local bend. (c) hand specimen. (d) Scanning electron microscopy (SEM) images. Detailed petrographic examination by SEM and transmitted light showing mineral assemblage: (e) Siderite. (f) Hematite. (g) Chlorite. (h) Sericite. Cht = Chert, Mt = Magnetite, Sd = Siderite, Q = Quartz, Hem = Hematite, Chl = Chlorite, Ser = Sericite.




Fig. 4 Concordia diagram of the zircon U-Pb ages of the tuff at the Baijianshan BIF.




Fig. 5 (a) Chondrite-normalized REE pattern of the basalts. Chondrite-normalized values after Sun and McDonough (1989). (b) PAAS-normalized REE+Y diagrams of BIFs from the Baijianshan BIF. The grayish zone represents the REY data from the Xinyu BIF (Li et al., 2014). PAAS-normalized values after McLennan (1989).




Fig. 6 (a) Bi-variation between Rb and Th of Baijianshan BIF samples; (b-i) Bi-variation between Al_2O_3 and TiO_2 (b), K_2O (c), Rb (d), V (e), Sc (f), Cs (g), Cr (h) and REE (i) of the Baijianshan BIF. Data for bulk-rock BIF samples are listed in supplementary Table 2.




Fig. 7 $^{87}\text{Sr}/^{86}\text{Sr}_i$ diagram of BIF and basalt samples in Baijianshan.




Fig. 8 ϵ_{Nd} ($t = 0.737\text{Ga}$) vs. Al_2O_3 diagram of the Baijianshan BIF; (b) $\text{Ce}/\text{Ce}^*_{\text{PAAS}}$ vs. Al_2O_3 diagram of the Baijianshan BIF.




Fig. 9 Sm/Yb vs. Eu/Sm ratios illustrating two component mixing model of the hydrothermal fluid of the Baijianshan BIF (after Alexander et al., 2008). BJS-Baijianshan BIF; XY-Xinyu BIF (Li et al., 2014); DHLT-Dahongliutan BIF (Hu et al., 2017). Data sets of Archean-Paleoproterozoic BIFs are from Planavsky et al. (2010). Average compositions of high-T ($> 300\text{ }^\circ\text{C}$) hydrothermal fluids, low-T ($< 200\text{ }^\circ\text{C}$) hydrothermal solutions and Pacific seawaters were after Bau and Dulski (1999), Michard et al. (1983) and Alibo and Nozaki (1999), respectively.




Fig. 10 Pr/Pr^*_{PAAS} vs. Ce/Ce^*_{PAAS} diagram for the BIF at Baijianshan (after Bau and Dulski, 1996). The light blue area represents the data of Archean-Paleoproterozoic BIFs. Data sets of other BIFs are same as those of Fig. 9.




Fig. 11 $\delta^{57}Fe_{IRMM-014}$ of various ages BIFs in the world (Data of SW Greenland BIFs are from Dauphas et al.(2004); Manjeri Formation, Belingwe Belt, Zimbabwe BIFs are from Rouxel et al. (2005); Anshan-Benxi BIFs are from Li et al. (2012); Transvaal BIFs are from Johnson et al. (2003); Yuanjiaocun BIFs are from Hou et al. (2014); Gongchangling BIFs from Li et al. (2014); Sanjiang BIF are from Yan et al. (2010) and Xinyu BIF are from Shen et al. (2008)).




Fig. 12 Cartoon model of the formation of the Baijianshan BIF (see details in the text).

Appendix Table Captions

Appendix-Table 1 LA-ICP-MS U-Pb data for zircon from tuff in the first section of the Lapeiquan Formation

Appendix-Table 2 Geochemical compositions of the Baijianshan BIF

Appendix-Table 3 Sr-Nd isotopic data of the Baijianshan BIF

Appendix-Table 4 Fe isotopic data of the Baijianshan BIF

Supplementary Table 1 Zircon U-Pb age data of the tuff from the Baijianshan BIF

Spot	U ppm	Th ppm	Th/U	²⁰⁶ Pb/ ²³⁸ U Age	²⁰⁷ Pb/ ²³⁵ U Age	²⁰⁷ Pb/ ²⁰⁶ Pb	²⁰⁷ Pb/ ²³⁵ U	²⁰⁶ Pb/ ²³⁸ U	²⁰⁶ Pb/ ²³⁵ U	²⁰⁶ Pb/ ²³⁸ U	²⁰⁶ Pb/ ²³⁸ U		
2071-1	3079	831	0.27	1335	14	1360	21	0.0887	0.0012	2.8156	0.0427	0.2301	0.0025
2071-2	805	427	0.53	898	11	1154	20	0.1027	0.0014	2.1169	0.0369	0.1495	0.0018
2071-3	529	381	0.72	776	8	1009	18	0.0965	0.0016	1.7019	0.0307	0.1279	0.0014
2071-4	155	81	0.52	835	9	1350	22	0.1458	0.0022	2.778	0.0462	0.1382	0.0015
2071-5	184	101	0.55	751	8	1009	18	0.0999	0.0016	1.7022	0.0304	0.1236	0.0013
2071-6	297	252	0.85	1273	14								

Supplementary Table 2 Geochemical compositions of Baijianshan BIF

Sample	BJS-1	BJS-2	BJS-3	BJS-4	BJS-5	BJS-6	BJS-7	BJS-8	BJS-9	BJS-10
Major elements (%)										
Fe ₂ O ₃	53.60	41.52	56.64	52.26	30.01	36.85	41.01	45.22	39.78	48.55
SiO ₂	29.91	38.96	26.81	31.20	40.92	40.79	36.97	41.77	43.66	37.59
Al ₂ O ₃	1.56									

TFe ₂ O ₃	21.79	28.86	44.82	42.63	26.98	13.11	40.45	11.61	11.70	11.98
SiO ₂	55.06	49.87	41.24	43.65	53.64	59.90	42.80	39.94	40.91	44.55
Al ₂ O ₃	5.24	4.93	2.81	2.86	4.83	7.58	3.84	11.36	11.45	10.76
CaO	0.81	1.02	0.75	0.75	0.76	0.84	0.65	9.77	9.31	7.45
MgO	1.57	1.51	0.74	0.86	1.36	2.32	1.31	8.40	8.54	10.38
TiO ₂	0.60	0.53	0.35	0.25	0.59	0.86	0.41	3.66	3.80	3.60
MnO	0.03	0.04	0.05	0.03	0.03	0.02	0.03	0.24	0.25	0.23
P ₂ O ₅	0.20	0.35	0.49	0.40	0.23	0.19	0.54	0.46	0.43	0.38
K ₂ O	3.11	2.32	0.26	0.17	1.86	3.78	1.98	1.35	1.27	0.36
Na ₂ O	0.94	1.25	0.14	0.19	0.63	0.77	0.88	0.94	0.98	0.59
LOI	2.06	1.97	4.57	2.89	2.9	3.19	1.92	12.02	11.39	9.60
FeO	7.97	7.26	3.89	5.29	6.06	6.57	5.31	-	-	-
Total	99.38	99.91	100.11	99.97	99.87	99.13	100.12	99.75	100.03	99.88
Trace elements (ppm)										
Cr	118	81.0	59.7	10.5	86.2	112	50.9	214	214	306
Ni	15.6	14.5	12.1	11.0	15.6	12.3	12.6	162	166	217
Co	76.1	70.2	33.7	34.7	54.0	27.1	45.3	48.0	62.9	50.7
Sc	12.2	14.4	7.97	8.11	11.7	15.1	8.11	26.4	26.1	25.7
Ba	662	562	111	19.2	359	1219	385	877	864	296
Sr	45.0	53.3	45.8	26.1	32.6	37.0	48.9	205	209	150
Rb	111	95.6	7.36	6.13	113	184	85.6	34.4	32.2	12.3
Zr	101	93.4	60.0	87.9	79.9	116	79.7	259	251	228
Hf	2.63	2.37	1.23	1.50	2.11	2.99	1.92	5.35	5.74	6.00
Ta	0.88	0.75	0.37	0.36	0.64	0.84	0.53	2.24	2.33	2.14
Ga	7.95	7.63	4.39	4.63	7.12	10.3	5.97	18.3	17.2	18.8
Cu	27.6	21.6	8.12	3.56	21.4	36.0	12.4	-	-	-
Zn	62.1	40.0	26.0	23.8	35.3	37.0	29.6	-	-	-
Pb	7.80	8.24	3.54	5.32	7.64	15.4	5.75	-	-	-
U	0.47	0.69	0.39	0.40	0.72	0.82	0.37	1.45	1.09	0.92
Th	7.26	5.14	2.81	2.97	5.52	7.11	4.11	3.32	3.37	3.18
Nb	8.35	8.21	4.19	4.86	6.80	9.50	6.11	38.0	36.5	33.6
La	26.8	20.3	13.0	9.82	19.1	22.5	17.3	36.9	34.1	33.7
Ce	58.3	44.2	30.2	22.8	42.0	46.1	39.0	77.2	72.9	74.9
Pr	7.03	5.35	3.73	2.77	5.09	5.51	4.81	9.37	9.15	9.28
Nd	29.3	22.4	16.2	11.7	21.3	22.6	20.7	37.9	36.6	40.0
Sm	6.46	4.95	3.67	2.65	4.70	4.96	4.70	8.06	8.20	9.11
Eu	1.33	1.15	0.91	0.66	1.06	1.01	1.12	1.41	1.85	2.91
Gd	6.39	5.03	4.02	3.03	4.64	4.93	5.01	7.69	7.82	8.87
Tb	1.00	0.80	0.67	0.52	0.72	0.79	0.82	1.21	1.22	1.35
Dy	5.98	4.98	4.53	3.71	4.26	4.84	5.36	6.57	7.04	7.04
Y	28.5	24.9	27.9	22.0	20.1	26.1	31.9	34.8	34.0	32.1
Ho	1.20	1.04	1.02	0.85	0.86	1.01	1.17	1.25	1.29	1.31
Er	3.23	2.94	3.00	2.59	2.33	2.82	3.38	3.11	3.15	3.20
Tm	0.45	0.43	0.46	0.40	0.34	0.41	0.50	0.39	0.40	0.39
Yb	2.87	2.93	3.06	2.76	2.20	2.69	3.36	2.28	2.36	2.28
Lu	0.43	0.45	0.49	0.45	0.33	0.41	0.53	0.34	0.32	0.32
	150.77	116.95	84.95	64.71	108.93	120.58	107.75	193.68	186.4	194.67
Y/Ho	23.75	23.94	27.35	25.76	23.29	25.84	27.26	27.84	26.36	24.50
La/La*	1.06	1.07	1.09	1.02	1.06	1.09	1.10	1.14	1.06	1.14
Ce/Ce*	0.98	0.98	0.99	1.00	0.98	0.95	0.98	0.98	0.98	1.01
Y/Y*	0.85	0.88	1.04	0.99	0.84	0.95	1.02	0.89	0.83	0.78
Pr/Pr*	1.00	1.00	0.99	0.99	1.00	1.00	0.99	0.97	0.99	0.95
Eu/Eu*	0.97	1.08	1.10	1.08	1.07	0.96	1.08	0.54	0.69	0.97
La _N /Yb _N	0.69	0.51	0.31	0.26	0.64	0.62	0.38	10.99	9.82	10.04

Supplementary Table 3 Sr-Nd isotopic data of the Baijianshan BIF

Sample	⁸⁷ Rb/ ⁸⁶ Sr	⁸⁷ Sr/ ⁸⁶ Sr	(⁸⁷ Sr/ ⁸⁶ Sr) _i	¹⁴⁷ Sm/ ¹⁴⁴ Nd	¹⁴³ Nd/ ¹⁴⁴ Nd	T _{DM} (Ga)			
BJS-1	0.0433	0.717927	0.000004	0.7175	0.1457	0.512425	0.000006	1.63	0.65
BJS-4	4.2608	0.727751	0.000009	0.6829	0.1308	0.512394	0.000003	1.39	1.46
BJS-5	1.4305	0.728893	0.000008	0.7138	0.1378	0.512363	0.000003	1.58	0.19
BJS-7	2.4192	0.729423	0.000009	0.7040	0.1473	0.512376	0.000002	1.78	-0.45
BJS-8	2.8555	0.718681	0.000003	0.6886	0.1238	0.512351	0.000004	1.36	1.27
BJS-10	2.6390	0.723954	0.000005	0.6962	0.1344	0.512364	0.000002	1.51	0.53
BJS-12	7.0891	0.740795	0.000007	0.6662	0.1333	0.512311	0.000003	1.59	-0.40
BJS-16	9.9936	0.746329	0.000008	0.6412	0.1334	0.512317	0.000003	1.58	-0.30
BJS-18	5.0054	0.731954	0.000008	0.6793	0.1373	0.51237	0.000004	1.56	0.38
2702H1	0.4737	0.711034	0.000009	0.7061	0.1286	0.512662	0.000003	0.88	6.90
2702H2	0.4355	0.711156	0.000009	0.7066	0.1354	0.512694	0.000006	0.89	6.88
2702H3	0.2322	0.710822	0.000012	0.7084	0.1377	0.512644	0.000005	1.02	5.69

Supplementary Table 4 Fe isotopic data of the Baijianshan BIF

Sample	⁵⁷ Fe _{IRMM-014}	⁵⁶ Fe _{IRMM-014}
--------	--------------------------------------	--------------------------------------

BJS-2	2.10	0.09	1.46	1.46
BJS-5	1.78	0.02	1.24	1.24
BJS-7	2.76	0.09	1.89	1.89
BJS-9	2.49	0.05	1.69	1.69
BJS-12	3.05	0.04	2.17	2.17
BJS-17	2.80	0.01	1.91	1.91
